

Routine childhood immunisation programme

Each vaccination is given as a single injection into the muscle of the thigh or upper arm.

At what age to immunise	To protect against	Vaccine given
Two months old	Diphtheria, tetanus, pertussis (whooping cough), polio and <i>Haemophilus influenzae</i> type b (Hib) Pneumococcal disease	DTaP/IPV/Hib and Pneumococcal conjugate vaccine (PCV)
Three months old	Diphtheria, tetanus, pertussis, polio and Hib Meningococcal group C disease (MenC)	DTaP/IPV/Hib and MenC
Four months old	Diphtheria, tetanus, pertussis, polio and Hib MenC Pneumococcal disease	DTaP/IPV/Hib and MenC and PCV
Between 12 and 13 months old – within a month of the first birthday	Hib MenC Pneumococcal disease Measles, mumps and rubella (German measles)	Hib/MenC, PCV and MMR
Three years and four months or soon after	Diphtheria, tetanus, pertussis and polio Measles, mumps and rubella	DTaP/IPV or dTaP/IPV and MMR
Girls aged 12 to 13 years	Cervical cancer caused by human papillomavirus types 16 and 18	HPV
13 to 18 years old	Tetanus, diphtheria and polio	Td/IPV

My child's next appointment:

Questions?

Ask your practice nurse, health visitor
or call Tracy Fabiano, Immunisation
Co-ordinator, on 01392 267 667
or text 07837 685 347.

Or visit: <http://www.gosh.nhs.uk/immunisation/faqs.html> or
<http://www.nhs.uk/Planners/vaccinations/Pages/Aboutvaccinationhub.aspx>



Don't
Hesitate -
Vaccinate 🚫